iPLEDGE-Committed to Pregnancy Prevention P.O. Box 29094 Phoenix, AZ 85038

Start to feel sad or have crying spells

Before I start taking isotretinoin, I agree to tell my doctor if, to the best of my knowledge, anyone in my family has ever had symptoms of depression, been psychotic, attempted suicide, or had any other serious mental problems.

Once I start taking isotretinoin, I agree to stop using isotretinoin and tell my doctor right away if any of the following signs and symptoms of depression or psychosis

Patient Information/Informed Consent (for all patients):

To be completed by patient (and parent or guardian if patient is under age 18) and signed by the doctor.

Read each item below and initial in the space provided if you understand each item and agree to follow your doctor's instructions. A parent or guardian of a patient under age 18 must also read and understand each item before signing the agreement.

Do not sign this agreement and do not take isotretinoin if there is anything that you do

Do not sign this agreement and do not take isotretinoin if there is anything that you do not understand about all the information you have received about using isotretinoin			Lose interest in activities I once enjoyed	
110	t didenstand about an the information you have received about using isotretinon.		Sleep too much or have trouble sleeping	
1.	I,(Patient's Name)		Become more irritable, angry, or aggressive than usual (for example, temper outbursts, thoughts of violence)	
	understand that isotretinoin is a medicine used to treat severe nodular acne that cannot be cleared up by any other acne treatments, including antibiotics. In severe nodular acne, many red, swollen, tender lumps form in the skin. If untreated, severe nodular acne can lead to permanent scars.		Have a change in my appetite or body weight	
			Have trouble concentrating	
			Withdraw from my friends or family	
	Initials:		Feel like I have no energy	
2	My doctor has told me about my choices for treating my acne.		Have feelings of worthlessness or guilt	
2.	Initials:		Start having thoughts about hurting myself or taking my own life (suicidal thoughts)	
3.	I understand that there are serious side effects that may happen while I am taking		Start acting on dangerous impulses	
	I understand that there are serious side effects that may happen while I am taking isotretinoin. These have been explained to me. These side effects include serious birth defects in babies of pregnant patients. [Note: There is a second Patient Information/Informed Consent About Birth Defects (for female patients who can get pregnant)].		Start seeing or hearing things that are not real	
			Initials:	
	Initials:		agree to return to see my doctor every month I take isotretinoin to get a new prescription for isotretinoin, to check my progress, and to check for signs of	
4.	I understand that some patients, while taking isotretinoin or soon after stopping isotretinoin, have become depressed or developed other serious mental problems. Symptoms of depression include sad, "anxious" or empty mood, irritability, acting on dangerous impulses, anger, loss of pleasure or interest in social or sports activities, sleeping too much or too little, changes in weight or appetite, school or work performance going down, or trouble concentrating. Some patients taking isotretinoin have had thoughts about hurting themselves or putting an end to their own lives (suicidal thoughts). Some people tried to end their own lives. And some people have ended their own lives. There were reports that some of these people did not appear depressed. There have been reports of patients on isotretinoin becoming aggressive or violent. No one knows if isotretinoin caused these behaviors or if they would have happened even if the person did not take isotretinoin. Some people have had other signs of depression while taking isotretinoin (see #7).	8	ide effects.	
			Initials:	
		9. I	sotretinoin will be prescribed just for me $-$ I will not share isotretinoin with the people because it may cause serious side effects, including birth defects.	
			Initials:	
		10. I	will not give blood while taking isotretinoin or for 1 month after I stop taking sotretinoin. I understand that if someone who is pregnant gets my donated blood, er baby may be exposed to isotretinoin and may be born with serious birth defect	
			Initials:	
		11 I		
	Initials:	11. I	I have read <i>Patient Introductory Brochure</i> and other materials my provider provided me containing important safety information about isotretinoin. I understand all the information I received.	
	Before I start taking isotretinoin, I agree to tell my doctor if I have ever had symptoms of depression (see #7), been psychotic, attempted suicide, had any other mental problems, or take medicine for any of these problems. Being psychotic means having a loss of contact with reality, such as hearing voices or seeing things that are not there.	U	Initials:	
		10 N		
		12. b	My doctor and I have decided I should take isotretinoin. I understand that I must e qualified in the iPLEDGE Program to have my prescription filled each month. I nderstand that I can stop taking isotretinoin at any time. I agree to tell my doctor	
	Initials:	if	I stop taking isotretinoin.	
	Initials.		Initials:	
Ind	ow allow my doctor to	begin 1	ny treatment with isotretinoin.	
Patient Signature:		I	Date:	
Par	ent/Guardian Signature (if under age 18):		Date:	
Pat	ient Name (print)			
Pat	ient Address		Telephone	
r 1				
I ha				
			ture and purpose of isotretinoin treatment, including its benefits and risks	
tr	rovided the patient the appropriate educational materials, such as the <i>Patient Introductory Brochure</i> eatment with isotretinoin	e and asl	sed the patient if he/she has any questions regarding his/her	
o ai	nswered those questions to the best of my ability			
Do	ctor Signature:		Date:	
PL	ACE THE ORIGINAL SIGNED DOCUMENTS IN THE PATIENT	'S ME	EDICAL RECORD.	

PLEASE PROVIDE A COPY TO THE PATIENT.